
Cumulative Risk Assessment: Residential & Food Exposure Methodology for the Organophosphorus Pesticides

Technical Briefing



U.S. EPA
Office of Pesticide Programs

November 15, 2001





Welcome and Introduction

Lois Rossi, Director Special Review and Reregistration Division

1/19/2001 13:02



Goals of Today's Meeting

Lois Rossi

Director

Special Review & Reregistration Division



oals & Purpose of Briefing

- Part of process the Agency has developed in conjunction with CARAT Workgroup on OP Cumulative Process
 - Update stakeholders
 - Provide explanations of methods as early as possible



oal

- Stakeholders understand methods & data so that they can provide effective input
- Find ways of enhancing transparency and ease of stakeholder participation



umulative Assessment Represents ew "Paradigm"

- Clearly not just a compilation of individual chemical risks
- Different way of looking at risk
 - Different questions
 - Different methods
 - Different risk management considerations



eveloping Common Understanding f New Paradigm

- Natural to want to know the "bottom-line"
 - What are the risk numbers?
 - What will be the regulatory outcome?
 - Will these methods be used for other groups?



eveloping Common Understanding f New Paradigm

- Why "bottom-line" must wait
 - Numbers will not be fully QA/QCed until preliminary risk assessment is completed
 - Need to have a common understanding of the methods/paradigm before there can be meaningful dialog on regulatory outcome
 - Application to other groups will not be known immediately
 - Depends on conclusions regarding methodology
 - Depends on specific characteristics of the other groups



Inderstanding New Methods

- A big piece is analyzing variability
 - What are the major factors contributing to risk
 - i.e. what matters & what does not
 - Analyze information sources
 - Refine risks where necessary
 - Understanding uncertainty
 - Agency committed to doing this



n Summary

- Focus on methods
- Prepare for release of preliminary assessment
- Stakeholders prepared to review preliminary assessment
- Understand what is important in preliminary assessment
- Contribute to the Agency's ability to conduct appropriate refinements
- Prepare proper foundation for analyzing risk mitigation, if necessary



gency & Stakeholder Cooperation

- Due to development of many new methods for cumulative, the Agency is providing briefings on these methods & the status of the assessment before the preliminary risk assessment is issued
- Stakeholders must recognize certain elements (both small & potentially large) may change
- Therefore, we all must work together cooperatively; to use the information constructively; and for the purpose of providing useful input



teps for Conducting a umulative Risk Assessment

Identify Common Mechanism Group

Identify Potential Exposures

Characterize + Select Common Mechanism Endpoint(s)

Determine Cumulative Assessment Group

Determine Absolute & Relative Toxic Potencies of Chemicals & Point of Departure for Index Chemical

Develop Detailed Exposure Scenarios for all Routes & Durations

Establish Exposure Input Parameters

Conduct Final Cumulative Risk Assessment

Conduct Characterization of Cumulative Risk



Overview of Foday's Briefing

Margaret Stasikov Director, Health Effects Division



oadmap

- Probabilistic methods for cumulative
- Proposed method for dietary
 - Residential uses
- Proposed methods for residential
- Summary & next steps



ood & Residential Team Members

- Elizabeth Doyle
- Jeff Evans
- David Hrdy
- Carol Lang

- David Miller
- Mohsen Sahafeyan
- William O. Smith
- Bill Wooge



Food and Residential Exposures and the Risk Assessment Process

David Miller HED



oadmap

- Probabilistic methods for cumulative
- Proposed method for food
- Residential uses
 - Proposed methods for residential
- Summary & next steps



ntroduction

- Both food and residential exposure estimates used in a probabilistic assessment for cumulative risk
 - Presentation in October covered exposures through drinking water
- Next slides detail:
 - How food risk is estimated probabilistically
 - How residential exposures would need to be estimated to be combined in a probabilistic estimate with food
 - How food and residential exposures are combined



utline

- Risk Equation
- Using DEEM[™] software for assessments for food (alone)
- Key Concepts in Aggregation/Cumulation Methodology using DEEMTM /CalendexTM
 - Importance of calendar-based Assessment
- Illustrative step-by-step example of Probabilistic Aggregate/Cumulative Assessment for Food and Residential Exposures



isk Equation

- Risk is the probability that a substance will produce harm under specified conditions
- Risk = Exposure X Hazard
 - Hazard part derived from toxicological studies
 - Exposure part derived from
 - FOOD: residues and consumption
 - RESIDENTIAL: residues and contact



xposure From Food

- We use DEEM[™] (Dietary Exposure Evaluation Model) software to estimate exposures from food
 - Exposure = Consumption X Residue
 - DEEM[™] produces an estimate of the distribution of exposures through food
 - Uses probabilistic (Monte-Carlo) techniques



EEMTM Software

- Uses data on food consumption and data on pesticide concentration to estimate exposures from food
 - Food Consumption
 - USDA's Continuing Survey of Food Intakes by Individuals (CSFII)
 - Survey reports daily consumption of food for 40,000+ person-days of consumption
 - 1994-96/1998 data available
 - Pesticide Residues
 - Market Basket Data
 - USDA PDP data
 - FDA Monitoring data




ey Concepts in Cumulative Assessment opropriate Matching and Combining

Objective: to appropriately match and subsequently combine estimates of pesticide exposures through food with estimates of pesticide exposures through residential uses

Key Concepts in Cumulative Assessment Appropriate Matching and Combining

- Matching and combining must appropriately consider temporal and spatial factors associated with exposure
 - Temporal Factors
 - The time of year that pesticide exposures occur
 - E.g., springtime
 - Pesticide exposures on one day can be related to pesticide exposures on previous day
 - E.g. day-to-day relationships
 - Spatial Factors
 - Region of Country in which pesticide exposures occur
 - E.g., South vs. North

Regional Framework



Source: USDA ERS



DEEMTM/CalendexTM Cumulative Assessmen

DEEM™/Calendex™ provides a probabilistic assessment in which appropriate matching occurs

- Incorporates concept of a Calendar to evaluate aggregate exposures
- Looks at each individual day of the year
 - Allows appropriate "temporal matching" of exposures through food, drinking water, and residential pathways.
 - Temporal aspect of exposure through residential uses important for OP's due to expected seasonal use-patterns



EEM™/Calendex Cumulative Assessme

- What would happen if we didn't use calendarbased approach?
- For example:
 - Fall dermal exposure through lawn-use could be (incorrectly) combined with dermal exposure through spring flea treatment on pets
 - Oral hand-to-mouth exposure from spring lawn application on one day could be (incorrectly) combined with drinking water concentration characteristic of the winter season



ey Concepts in Cumulative Assessment opropriate Matching and Combining

In summary, must track potentially exposed persons on a daily basis in a way that preserves all appropriate linkages in a way that considers time, region, and age groups



- For each assessment region:
 - Maintain temporal & spatial characteristics
 - Use DEEM™ software to estimate exposure from food
 - Use Calendex[™] software to aggregate/cumulate exposure from food and residential exposures and drinking water



ge Groups

- Assessment performed for the following age groups:
 - Children 1-2 years old
 - Children 3-5 years old
 - Adults 20-49 years old
 - Adults 50+



lustrative Example of Calendex $^{ extsf{TM}}$ Analy

- 1-day exposure is presented as an example
- Analysis serves as "building block" for any number of days analysis
- Only food & residential included in this example



lustrative Example of Calendex $^{ extsf{TM}}$ Analy

Hypothetical Consumption Profile for CSFII Individual #

12 kg child

Consumed: 260 g food #1

320 g food #2

250 g food #3

- Period of Interest: January 1 through December 31
- Specific to Region of Interest



ustrative Example of Calendex™ Analysis

- STEP 1: Calculate Exposure from Food for Individual # on January 1
 - Food Exposure(from DEEM™): = 2.89x 10⁻⁵ mg/kg bw/day
- STEP 2: Select Residential Treatments for Individual # on January 1
 - Specific to region & time and demographics of individual
 - Assigned probabilistically
 - Were pesticides applied in/around home?
 - If so, which treatments?
 - And how much, how often, during what time frame, with what frequency, and by whom?

Ilustrative Example of CalendexTM Analysis

- STEP 3: Calculate Exposure from any assigned new residential uses for January 1
- STEP 4: Determine if Exposure is "Active" from any previously assigned use/application
 - by oral (hand to mouth) exposure to children (2 days earlier)
- Exposure = $\{ISR 2DAT \times Surf Area_{finger} \times No. events/hr \times No. hours \times Saliva Removal efficiency\}$
- body weight
 - $= 1.33 \times 10^{-5} \text{ mg/kg BW/day}$
 - STEP 5: Aggregate exposures for Day #1 from Food and (any active) Residential Uses
- = 2.89x 10⁻⁵ mg/kg BW/day + 1.33 x 10⁻⁵ mg/kg BW/day



Illustrative Example of CalendexTM Analysis

- STEP 6: Repeat Steps 1-5 many additional times for this individual, randomly selecting a series of treatment scenarios for that year, determining if any are applicable or otherwise "active for Day #1 for that individual, and aggregating (summing) selected food and residential exposures
- STEP 7: Continue steps 1-6 with Individual #2 through Individual # ~20,000
 - Result is a collection (or distribution) of aggregate exposures for January 1 for the relevant region
- STEP 8: Repeat steps 1-7 for January 2
 - Result is a collection (or distribution) of aggregate exposures for January 2 for the relevant region




lustrative Example of Calende x^{TM} nalysis

- STEP 9: Repeat steps 1-8 for January 3 through December 31
 - Result is a collection (or distribution) of aggregate exposures (food and residential combined) for each day of the year for the relevant region
 - These exposures can be plotted as a "time-line" or profile of daily exposures for any given percentile in the distribution



Example of CalendexTM Analysis (time based exposure profile)



Day of the Year



ummary

- Food and residential exposures and water will be considered probabilistically in the cumulative assessment
 - Reflects realistic pesticide use based on pest pressures, weather, activity patterns, etc.
 - Temporal and spatial characteristics will be preserved and maintained to produce realistic assessments



ummary

- Result of Assessment will be a time based exposure profile of exposures at any selected percentile
 - Total Exposure and its food, residential (and water) components
 - Single-day assessments will serve as "building block"



COMULATIVE DIETARY EXPOSURE FROM USES OF ORGANOPHOSPHORUS PESTICIDES

William O. Smith Health Effects Division



urpose

Describe methods used and assumptions involved in cumulative dietary exposure assessment for organophosphorus pesticides using an RPF approach and the DEEM™ model



utline of Presentation

- Scope of food exposure assessment
- Approach to combining residues
- Calculation of residue inputs
- Discussion of assumptions in assessment



cope of Food Assessment

- Sources of Residue Data
- OPs included
- Foods included



cope of Food Assessment ources of Residue Data

- USDA Pesticide Data Program (PDP)
 http://www.ams.usda.gov/science/pdp
- FDA Center for Food Safety & Applied Nutrition

http://www.cfsan.fda.gov/~lrd/pestadd.html
Pesticide Residue Monitoring Program
Total Diet Study (TDS)



USDA Pesticide Data Program

- Designed to provide data on dietary exposure
 - Started in 1991
 - Samples foods highly consumed by children
 - Reflect foods typically available throughout year
 - Foods collected near point of consumption
 - Food washed and inedible portions removed before analysis



DA Monitoring Data

- Designed to enforce EPA tolerances in imported foods and in domestic foods shipped in interstate commerce
- Domestic samples are collected as close as possible to the point of production in the distribution system
- Import samples are collected at the point of entry into U.S. commerce
- Emphasis is on the raw agricultural product, which is analyzed as the unwashed, whole (unpeeled), raw commodity. Processed foods are also included



DA Total Diet Study

- Provided data on dietary intake of food contaminants for almost 40 years
- Since 1991 26 market baskets collected
 - Each MB includes ~ 260 foods collected in grocery stores
 - Analyses performed on foods prepared for consumption
 - Highly sensitive analyses
 - Limited number of samples



Ps Included in Current Food Assessment

 Acephate, azinphos methyl, chlorpyrifos, chlorpyrifos-methyl, disulfoton, diazinon, dichlorvos, dimethoate, ethoprop, fenamiphos, malathion, methidathion, methamidophos, mevinphos, oxydemeton-methyl, methyl parathion, phorate, phosolone, phosmet, pirimiphos methyl, terbufos, & tribufos



Ps Not Included in Current Assessment

- OP uses that have been voluntarily cancelled
- OPs that only have public health uses
- OPs with no detectable residues in monitoring



DP Foods Included

oples ople Juice

ananas

occoli

elery

antaloupe

arrots

veet Corn

ıcumbers

orn Syrup

nerries

ce

Green Beans

Grapes

Grape Juice

Lettuce

Milk

Oats

Oranges

Orange Juice

Peaches

Pears

Nectarines

Pineapple

Potatoes

Bell Peppers

Strawberries

Sweet Potatoes

Soybean

Spinach

Sweet Peas

Tomatoes

Wheat

Winter Squash

Poultry

Peanut Butter



DP Food Types

ruits and Vegetables **Grains**

Whole Fresh

Frozen

Canned Milk

Whole

ruit Juices

Ready-to-Drink

Frozen

Concentrate

Poultry

Fat

Liver

Muscle



DP Food Coverage

- 44 Food Types Representing 86% of the Diet of Children 3-5
 - (CSFII 94-96,98)



oods Based on Translated PDP Data

Eggplant

Pepper(other than green)

Citrus (other than orange)

Apricots

Plums/Prunes

Rye

Beets-garden

Horseradish

Parsnips

Radishes

Rutabagas

Turnips

Various leafy greens

Brussels sprouts

Cabbage

Cauliflower

Melons(other than

cantaloupe)

Pumpkins

Squash-summer

Represent 1.3% of Diet of Children 3-5 (CSFII)



oods Based on FDA Monitoring Data

- Eggs
 - Assume negligible based on FDA monitoring data
- Seafood
 - Assume negligible based on FDA monitoring data
- Meat from Beef, Pork, Sheep & Goats
 - Used maximum residues found in FDA/TDS

Represent 6.3% of Diet of Children 3-5 (CSFII)



oods Assumed Negligible

- Sugarcane, Sugar Beet & Maple
 - Molasses, syrup & sugar
 - Assume negligible residues
 - Highly processed/refined
 - No residues in sugar or pancake syrup analyzed by FDA/TDS
 - No residues in corn syrup analyzed by PDP

Represent 3.1% of Diet of Children 3-5 (CSFII)



otal Food Forms Included in Assessment

- DEEM: 547 foods based on CSFII
- OP food assessment: 258 foods
 - 154 expanded from 44 PDP commodities
 - 54 by translation of PDP data
 - 29 meats from TDS data
 - 12 fish and eggs from FDA
 - 9 sugar, molasses, & syrup by inference



od Forms Not Included in Current Assessment

- Highly processed foods
- Infrequent detectable residues
- Residues detected negligible



The Proportion of the Diet of Children (3-5) Covered in the Cumulative assessment

Source of Residue Estimate	% of Diet		
PDP	85.7		
Translation of PDP	1.3		
FDA	6.3		
Assumed negligible	3.1		
Not included	3.6		




utline of Presentation

- Scope of food exposure assessment
- Approach to combining residues
- Calculation of residue inputs
- Discussion of assumptions in assessment



PF Approach

Converts chemical specific residues (B,D,P,T) on a food sample to a common residue (T)




wo Steps Needed to Derive umulative Residue_{ie}

1. $Residue_{IE} = Residue X PF X RPF$

PF=processing factor


2. Cumulative Residue_{IE} = Residue_{IE} (per PDP sample)





tep #1: Index Equivalent esidue (Residue_{IE})

 $Residue_{IE} = Residue X PF X RPF$





alculation Parameters for Residue_{IE}


 $Residue_{TF} = Residue X PF X RPF$

- Residue: PDP residue data by sample
- PF: Processing factors from single chemical assessments.
- **RPF:** Relative Potency Measure.



tep 2: Cumulative Residue

Cumulative Residue_{IE} = Residue_{IE} (per PDP sample)






utline of Presentation

- Scope of food exposure assessment
- Approach to combining residues
- Calculation of residue inputs
- Discussion of assumptions in assessment




ata Tables (screen print)





DP Data Table (screen print)





rocessing Factor Table

Microsoft Access - [Master-FFT : Table]								
Elle Edit View Insert Format Records Tools Window Help								
CropGroup	Commodity	Foodform	Code	PF	PestCode	Exclud		
0	Grapes	Uncooked	1311	1	P			
0	Grapes	Uncooked	1311	1	T			
0	Grapes	Uncooked	1311	1	M			
0	Grapes	Uncooked	1311	1	J			
0	Grapes	Uncooked	1311	1	S			
0	Grapes	Uncooked	1311	1	W			
0	Grapes	Uncooked	1311	1	V			
0	Grapes	Uncooked	1311	1	A			
0	Grapes	Uncooked	1311	1	R			
0	Grapes	Uncooked	1311	1	RA			
0	Grapes	Uncooked	1311	1	С			
0	Grapes	Uncooked	1311	1	В			
0	Grapes	Uncooked	1311	1	FB			
	Insert Format CropGroup CropGroup C C C C C C C C C C C C C	CropGroup Commodity CropGroup Commodity Crapes Crapes	CropGroup Commodity Foodform O Grapes Uncooked O Grapes Uncooked	CropGroup Commodity Foodform Code O Grapes Uncooked 1311 O Grapes Uncooked 1311	CropGroup Commodity Foodform Code PF Grapes Uncooked 1311 1 Grapes Uncooked 1311 1	CropGroup Commodity Foodform Code PF PestCode O Grapes Uncooked 1311 1 P O Grapes Uncooked 1311 1 M O Grapes Uncooked 1311 1 M O Grapes Uncooked 1311 1 S O Grapes Uncooked 1311 1 W O Grapes Uncooked 1311 1 R O Grapes Uncooked 1311 1 R		




elative Potency Factors Table

 Identifies Relative Potency Factors for all Chemicals in the assessment



Translation Table

- Links PDP foods and types to DEEM food form codes
 - Correlates PDP commodities with CSFII
 - Translations of data can be set in this table






umulative ResidueIE = Sum ResidueIE (per PDP sample)

icrosoft Access - [Query3-for-Picture : Select Query]

<u> Eile Edit View Insert Format Records Tools Window Help</u> 罗图マ 光 酯 🔒 💅 AL ZI 2 . **₽ Q** ♥ 4 ⊕ ね・ D# 10% CONCEN PF **RPF** Residue(IE SampID Chemical COMMOD FoodForm L9602140006GR Uncooked 0.55 0.07797 0.04288 R Grapes 1 L9602140006GR Uncooked 0.04274 P Grapes 0.44 0.09714 1 L9602140006GR W Grapes Uncooked 0 0.61022 0.00000 1 0 L9602140006GR V 0.00000 Grapes Uncooked 1 0.01661 L9602140006GR Τ Grapes Uncooked 0 1 1.00000 0.00000 L9602140006GR S 0.69178 Uncooked 0 1 0.00000 Grapes L9602140006GR R Grapes Uncooked 0 1 0.07797 0.00000L9602140006GR M Grapes Uncooked 0 1 0.09689 0.00000 L9602140006GR 0 Grapes Uncooked 1 0.30132 0.00000 J L9602140006GR C Grapes Uncooked 0 1 0.01090 0.00000L9602140006GR В Grapes Uncooked 0 1 0.01886 0.00000 L9602140006GR Grapes Uncooked 0.46537 0.00000 Α 0


Cumulative Residue Distribution





esidue Inputs for Food Assessment


- 220 Residue
 Distribution Files
 (RDFs) for different food forms
 - 201 point estimates for blended foods or other data sources





Cumulative Dietary Risk

Exposure = Residue X Consumption





onsumption

- USDA Continuing Survey of Food Intakes by Individuals
 - CSFII, 1994-1998



SFII 1994-96/1998

- Intakes of individuals residing in U.S.
 - 20,607 individual participants interviewed over two discontinuous days
- 1998 Supplemental Children's Survey
 - 5,559 additional children
 - Birth through 9 years old
 - Integrated into 1994-96 CSFII



SFII 1994-96/1998

The 1994-96/1998 CSFII significantly increases the number of children in the survey compared to the 1989-91 survey data currently being used by OPP

omparison of Number of 1-day and Complete Itakes, by Children's Age Group 1994-96/1998 vs. 1989-91 CSFII

ge	1994-96/1998 CSFII		1989-91 CSFII	
roup	No.	No.	No.	
	Individuals	Person-	Individuals	
		days		
1	1408	2	197	
2	2179	4114	574	
5	4579	8464	806	
11	2000	3706	1476	



onsumption

- USDA Continuing Survey of Food Intake by Individuals (CSFII) database
 - OP cumulative risk assessment uses 1994-96/1998 data with new (USDA/EPA) recipes
 - Respondents in the CSFII survey report what they ate in the form the food was eaten



Food to RAC Mapping/Translation

- Food consumption reported on an "as-eaten" basis
 - 200 g pizza
 - 100 g "Toasted Oat cereal"
- "Linked" or translated to the raw agricultural commodities for which we have residue data
 - Linking is done with USDA/USEPA recipes
 - Will be publicly available



onsumption

- Assessment based on survey information on the following age groups:
 - Children 1-2 years old
 - Children 3-5 years old
 - Adults 20-49 years old
 - Adults 50+ years old



lodel

- Dietary Exposure Evaluation Model DEEM-FCID™
- Probabilistic (Monte-Carlo) procedure
- Input:
 - Distributions for consumption
 - Distributions or point estimates for residue concentrations
- Output:
 - Distribution of one-day dietary exposures
 - Distribution of associated risks, i.e., MOEs



utline of Presentation

- Scope of food exposure assessment
- Approach to combining residues
- Calculation of residue inputs
- Discussion of choices & assumptions in assessment



- The assessment includes only chemical/crop combinations currently registered in U.S. or with import tolerances
 - Canceled & phased-out uses from single chemicals assessments are excluded
 - Violative residues are not included in assessment



- PDP samples were assumed to reflect residues as consumed
 - Used composite samples
 - Did not "decomposite" residues
 - PDP special surveys on single-servings or apples, pears, & peaches support use of composites



- PDP samples were assumed to reflect residues as consumed
 - Samples generally taken from wholesale distribution centers
 - Foods washed, peeled, etc. similar to preparation for consumption
 - Foods were not cooked



- Residue data in PDP are assumed to represent the potential for co-occurrence of OPs in single-day diets
 - Different foods consumed in one day may each contain OP residues
 - A single food may contain residues of more than one OP



- It was assumed that all OPs of concern were accounted for on each PDP sample
 - If an OP was not detected on a sample then it was assumed = zero
 - i.e., percent crop treated implicit in detection rate, subject to sensitivity of analytical method



- Higher percentiles of exposure not expected to be significantly affected by assumption of ND=0, given that cumulative residues significantly higher than LOD are abundant in the data
- This was demonstrated in the case study 12/99



- PDP residue data may be translated to foods not analyzed if similar agricultural practices are in effect
 - OPP/HED SOP 99.3



- PDP residue data may be translated to foods not analyzed if similar agricultural practices are in effect
 - 54 foods in this assessment were included by translation of PDP data
 - Translated to foods that are not major consumption items (~1% of child's diet)



- The food exposure component of the cumulative assessment is considered to be identical throughout the year and across regions, i.e., one national assessment will be performed.
- The national food estimate will be combined with 13 regional water assessments to provide for 13 regional dietary assessments.



uture Work

Impact of OP Market Basket Study



P Market Basket Study

- Currently being reviewed
 - 13 foods analyzed for 29 OP analytes
 - Single serving samples
 - Samples from grocery stores
 - Very low limits of detection
 - Good coverage of metabolites
- Preliminary analysis indicates similar dietary exposure estimate whether using PDP or Market Basket Survey.



P Market Basket Study

- Apples
- Broccoli
- Cherries
- Cucumbers
- Green beans
- Grapes
- Peaches

- Sweet corn
- Lettuce
- Orange
- Potatoes
- Strawberries
- Tomatoes



uture Work

- Analysis of exposure contributors
 - Pesticides
 - Foods
- Sensitivity analysis of omitted foods
- Incorporation of regulatory actions
- Incorporation of information from public comments



Current Status of OP Residential Uses

Deanna Scher

Special Review & Reregistration Division



7 OPs with Uses in Residential and/or ublic Areas Prior to Reassessment

- 6 OPs are now excluded entirely from the residential cumulative assessment
 - Diazinon
 - Dimethoate
 - Ethoprop
 - Fenitrothion
 - Phosmet
 - Propetamphos
- 11 OPs with remaining residential uses



esidential OP Usage: eduction of use inside the home

- Initially 9 OPs with indoor uses
 - Now only DDVP
 - Initially 6 OPs with pet uses
 - Now only tetrachlorvinphos and DDVP
 - Indoor use of chlorpyrifos, fenitrothion, and trichlorfon
 - Pre-packaged child-resistant bait stations (negligible exposure)



eduction in Overall OP Usage

- Diazinon + Chlorpyrifos
 - Residential use reduction of >20 million lbs. ai



rotecting Public Health Uses

- Public health uses have been retained where individual assessments indicate that these applications do not pose risks of concern and provide important public health benefits
 - Chlorpyrifos fire ant mound treatment
 - Chlorpyrifos mosquito control
 - Fenthion mosquito control
 - Naled mosquito and black fly control
 - Phosmet fire ant mound treatment



our OPs with Residential/Public Uses Have adividual Risk Mitigation

- Acephate, bensulide, chlorpyrifos, trichlorfon
- All remaining uses show risks below EPA's level of concern on an individual basis
 - The cumulative assessment reflects the most up-to-date residential use picture




our OPs with Residential/Public Ses Are Still Under Review

- Dichlorvos
- Fenamiphos
- Malathion
- Tetrachlorvinphos
- Any residential mitigation actions taken on these 4 OPs will be incorporated into the revised cumulative assessment



Residential Exposure Assessment Process

Jeff EvansHealth Effects Division



oadmap

- Probabilistic methods for cumulative
 - Proposed method for dietary
 - Residential uses
 - Proposed methods for residential
- Summary & next steps



urpose

- Use of a calendar based model to address the temporal use of residential OP's
 - Calendex™
- Use of distributions of data and exposure elements
- Use of survey data and other pesticide use information
- Approach is similar to the OP case study presented to SAP (12/7-8/00)



xpression of Residential Risk


- Routes considered, as appropriate
 - Oral, Dermal, Inhalation



cope

- Assessments conducted for 12 distinct geographical regions, reflecting climate & pest pressure differences
 - One region split into two residential assessments
- Includes remaining residential OPs that have significant exposure and exposure data
- Pet products not quantified
 - Only screening level SOPs available at this time

Regional Framework




Source: USDA ERS



egion 1 - Heartland

- Lawn use DDVP, Malathion, Trichlorfon
- Golf course Bensulide, Trichlorfon
- Ornamental gardens -Acephate, Disulfoton, Malathion
- Home gardens, etc. Malathion
- Indoor DDVP (pest strips and crack and crevice sprays)





egion 2 - Northern Crescent

Lawn - DDVP, Malathion, Trichlorfon


Golf course - Bensulide, Trichlorfon

Ornamental gardens - Acephate, Disulfoton, Malathion

Home gardens - Malathion

Indoor - DDVP (pest strips and crack and crevice treatments)

Public Health - Malathion, Naled





egion 3 – Northern Great Plains

- Lawn DDVP, Malathion, Trichlorfon
- Golf course Bensulide, Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice treatments)



YSNG, BS(HED, CEB)



egion 4 – Prairie Gateway

- Lawn Bensulide, DDVP, Malathion, Trichlorfon
- Golf course Acephate, Bensulide, Fenamiphos, Malathion, Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor (pest strips and crack and crevice treatments)
- Public Health Malathion





egion 5 – Eastern Uplands


- Lawn DDVP, Malathion, Trichlorfon
- Golf course Acephate, Bensulide, Fenamiphos, Malathion, Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home garden Malathion
- Indoor DDVP (pest strips and crack and crevice treatments)





egion 6 – Southern Seaboard

- Lawn Bensulide, DDVP, Malathion, Trichlorfon
- Golf course Acephate, Bensulide, Fenamiphos, Malathion, Trichlorfon
- Ornamental gardens -Acephate, Disulfoton, Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice treatments)
- Public health Malathion






egion 7a,b – Fruitful Rim, CA

(North Central and South Central Valley)

- Lawn DDVP, Malathion, Trichlorfon
- Golf course Bensulide, Fenamiphos, Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice treatments)





egion 8 – Basin & Range

- Lawn DDVP, Malathion, Trichlorfon
- Golf Course Bensulide, Trichlorfon
- Ornamental Gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice)

Region 8: Basin and Range





egion 9 – Mississippi Portal

- Lawn DDVP, Malathion, Trichlorfon
- Golf course Acephate,
 Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice)
- Public health Malathion, Fenthion






egion 10 – Fruitful Rim, NW

- Lawn, DDVP, Malathion, Trichlorfon
- Golf course none
 - Ornamental gardens Acephate, Disulfoton, Malathion
- Home gardens -Malathion
- Indoor DDVP (pest strips and crack and crevice)


Region 10: Fruit Rim, N





egion 11 – Fruitful Rim, TX


- Lawn Bensulide, DDVP, Malathion, Trichlorfon
- Golf courses Acephate,
 Bensulide, Fenamiphos,
 Malathion, Trichlorfon
- Ornamental gardens Acephate, Disulfoton,
 Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice)
- Public health Malathion





egion 12 – Fruitful Rim, FL

- Lawn DDVP, Malathion, Trichlorfon
 - Golf courses Acephate, Bensulide, Fenamiphos, Malathion
- Ornamental gardens Acephate, Disulfoton, Malathion
- Home gardens Malathion
- Indoor DDVP (pest strips and crack and crevice)
- Public health Fenthion, Malathion, Naled





cope/Registrations

- Lawn and Golf Course use
 - Lawn
 - Bensulide, dichlorvos, malathion, trichlorfon
 - Golf Course
 - Acephate, bensulide, fenamiphos, malathion, trichlorfon
- Home Garden
 - Acephate, disulfoton, malathion



cope/Registrations

- Indoor Crack and Crevice
 - Dichlorvos
 - **Pest Strips**
 - Dichlorvos
- Public Health
 - Individual assessments include concern
 - Naled, malathion, fenthion



cope/Routes - Lawns

- Trichlorfon, dichlorvos, malathion, bensulide (Texas only)
- Self applied, treatments made by adults
 - Exposure routes (Dermal and Inhalation)
- Post application exposure following hand and professionally applied treatments
 - Exposure routes Adults (Dermal)
 - Exposure routes Children (Dermal and Oral)



cope/Routes – Golf Course

- Acephate, trichlorfon, bensulide, fenamiphos, malathion
- Post application exposure following professionally applied treatments
 - Exposure Routes Adults (Dermal)



cope/Routes - Garden

- Garden
 - Shrubs, roses, flowers
 - Acephate, disulfoton, malathion
 - Applicator exposure (dermal, inhalation)
- Home crops
 - Malathion
 - Applicator exposure (dermal, inhalation)
 - Post application (dermal)



cope/Routes - Indoor

- Crack and crevice
 - Application and post application (inhalation)
- Pest strips
 - Post application (inhalation)
- Dichlorvos very volatile



cope - Public Health

- Wide area treatments (public health uses)
- Mosquitoes
 - Northern Crescent, Southern Seaboard, Mississippi Portal, Fruitful Rim (FL)
- Black Fly
 - Northern Crescent
- Post application
 - Dermal adults, children
 - Oral, children



awns – Use Information

- National Home & Garden Pesticide Use Survey (NHGPUS 1991)
- Doane Golftrak 1997, 1998
- Treated lawns based on regions using the National Garden Survey 1996-1997
 - Includes:
 - Percent of population participating in lawn care activities
 - Percent of population using insecticides
- Lawn Size (Vinlove and Torla 1995 and ORETF Survey)



awns – Use Information

- Chemical-specific turf residue data
- Certified/Commercial Pesticide Applicators
- State Cooperative Extension services
- Comparative Insecticide Effectiveness for Major Pest Insects of Turf in the United States



awn – Applicator Exposure Data

- Data source: ORETF
 - Application type:
 - Granular push-type rotary spreaders
 - Hose-end sprayer ready to use and one requiring the user to add the concentrate
- Clothing types:
 - Range of clothing
 - Short-sleeved shirt, short pants and long-sleeved shirt, long pants



awn – Post Application Exposure Data

- Residue transfer to skin (transfer coefficient)
 - Choreographed Activities of Adults Measured Using Biological Monitoring Vacarro 1996
 - Non-Scripted Activities of Children Measured Using Fluorescent Tracers Black 1993
- Chemical-specific lawn residue data



awn – Post Application Exposure Data

- Non-Dietary Ingestion (Hand-to-Mouth)
 - Hand-to-mouth frequency of events Reed 1999
 - Lawn residue data to account for saliva wetted hands Clothier 2000
 - Saliva extraction e.g., Camann 1995



olf Courses – Post Application Exposure Dat

- Percent of individuals participating in golf, 1992 Golf Course Operations by the Center for Golf Course Management
- Percent of Golf Courses Applying Selected Pesticides -Doane GolfTrak, 1998-1999.
- Residue transfer to skin (transfer coefficient)
 - Post application
 - Dermal route
- Chemical-specific turf residue data



ublic Health – Post Application

- Spray drift model
- Range of Deposition from applications based on use of AgDrift (as discussed in REDs)
- Deposition on lawns and post application is assessed in the same way lawn chemicals are
- Timing and pesticide used based on personal communication and publications prepared by organizations such as the Florida Coordinating Council of Mosquito Control



arden – Applicator Exposure Data

- Application of shaker can, garden duster, and small tank sprayer using proprietary data
- Applicator exposure dermal and inhalation
- Post application dermal exposure
 - Range of transfer coefficients from ARTF
- Duration of garden activities ORETF Survey




ndoor – Inhalation Exposure Data

- Dichlorvos range of inhalation exposure values for pressurized aerosol can (PHED)
- Post application inhalation exposure (adults and children)
- Pest strips
 - Collins et al., 1973
 - Crack and crevice
 - Gold et al., 1983
- Duration of time spent indoors
 - Exposure Factors Handbook



n Summary


- All available data considered
- Lawns
 - Lawn residue data available for all compounds
 - Regional adjustments where feasible
 - Home gardens DFR data from ARTF studies
 - CA and PA
- Wide Area treatments
 - Transfer efficiency based on chemical specific lawn residue data (except fenthion - assumed to be similar to malathion based on similar molecular weights and vapor pressure)





Summary & Next Steps

Lois Rossi Director Special Review & Reregistration Division



oadmap

- Probabilistic methods for cumulative
- Proposed method for food
- Residential uses
- Proposed methods for residential
- Summary & next steps



ummary of Cumulative Food & esidential Assessment Methods

- Food assessment:
 - Probabilistic assessment
 - Uses best available monitoring data
- Residential assessment:
 - Probabilistic assessment done with daily distribution of estimated residues
 - Regional assessment in order to combine residential (& water) on a realistic, localized scale
 - Focuses on realistic co-occurrences



ext Steps

 All data considered for the food & residential exposure assessment will be available with the preliminary assessment



ext Steps

November 2001

Release various revised science policy papers

December 2001

Release preliminary risk assessment

January 15, 2002

Technical briefing on preliminary cumulative risk assessment

January 16, 2002

CARAT Workgroup meets

Winter 2002 (Feb. 5-8 scheduled)

SAP Meeting on preliminary risk assessment

Winter 2002 (likely through March 8th)

Comment period on preliminary risk assessment

August 2002

Release revised risk assessment



ext Steps

For the most up-to-date information, visit:

http://www.epa.gov/pesticides/cumulative/

To receive OPP Updates via e-mail:

See the sign-up sheet at the registration desk